

A hybrid model for Rydberg gases including exact two-body correlations

Kilian P. Heeg,¹ Martin Gärttner,^{1,2} and Jörg Evers¹

¹*Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Deutschland*

²*Institut für Theoretische Physik, Ruprecht-Karls-Universität Heidelberg, Philosophenweg 16, 69120 Heidelberg, Germany*

(Dated: February 14, 2012)

A model for the simulation of laser-driven interacting Rydberg gases is discussed. Our hybrid approach combines an exact two-body treatment of nearby pairs of atoms with an effective treatment for separated pairs. A time-independent Monte Carlo technique is used to efficiently determine the steady state of the system. The hybrid model predicts features in the pair correlation function arising from multi-atom processes which existing models can only partially reproduce. Our interpretation of these features shows that higher-order correlations are relevant already at low densities.

PACS numbers: 32.80.Ee, 42.50.Nn, 67.85.-d

The distinctive properties of Rydberg atoms [1, 2] render them a powerful implementation of a tunable strongly interacting quantum many-body system. The signature effect of Rydberg atoms is the dipole blockade [3–8], which can be understood using rather basic theoretical models [8–10]. But ongoing experimental progress provides means to explore the correlations on a deeper level. Accordingly, more involved observables such as the Mandel Q parameter [11, 12], the pair correlation function [13] or quantum optical effects in the presence of Rydberg interactions [14, 15] moved into the focus of interest. State-of-the-art experiments push existing models for Rydberg gases to the limits of their validity ranges [13, 16], and require exceedingly long simulation times. Therefore, better simulation techniques are highly desirable.

One approach to simulate the many-body system is to truncate the otherwise exponentially growing state space, e.g., based on the Rydberg blockade. Such calculations of the exact Hamiltonian dynamics in a truncated Hilbert space provide valuable insights [9, 17–19]. But they are not well suited to model weakly interacting or low-density gases, as then the state space becomes too large. Due to the treatment on the level of the wave function, incoherent effects like spontaneous emission or dephasing cannot be incorporated in a straightforward way. Also, two-step excitation schemes cannot be described, as the intermediate state would spoil the state space reduction.

An alternative approach is to approximate the inter-atomic correlations. The simplest approach is the mean field approximation $\langle A_i B_j \rangle \approx \langle A_i \rangle \langle B_j \rangle$ for operators A, B acting on two different atoms i, j , which already describes the Rydberg blockade well [8, 10]. However, it fails for higher correlations and for more involved observables [14, 15]. A cluster expansion to higher-order atomic correlations could successfully describe an experiment on coherent population trapping in Rydberg atoms [14], but this method turned out to give inconsistent results at high densities [20]. Alternatively, a rate equation model was introduced in [21]. Using an adiabatic elimination of the atomic coherences, an effective rate equation for the atomic populations alone could be derived. The re-

sulting equation system still grows exponentially in the number of atoms, but can be solved using Monte Carlo techniques. This method is capable of modelling a multi-step excitation [20]. However, due to the simplified treatment of the interaction, only single-atom processes can be described and correlations are approximated.

In this work, we discuss a simulation technique combining higher predictive power with fast calculation times. In our model, atom pairs with distances below a characteristic length scale L_{Cr} are treated exactly, resulting in accurate two-body correlations. Atoms with distances above L_{Cr} are incorporated via effective detunings. We show that this hybrid approach describes three-body systems well over the whole range of interaction strengths. To efficiently determine the many-body steady state, we eliminate the time-consuming calculations of transition rates in each step based on a time-independent Monte Carlo technique. We show the consistency of our approach with the relevant existing models for simple observables as the excitation fraction of a disordered Rydberg ensemble. In contrast, the hybrid model predicts structures in the pair correlation function, which the existing models cannot fully reproduce. These structures




FIG. 1. (Color online) (a) Our model divides atoms into pairs and single atoms allowing an exact treatment of the two-body interaction up to a inter-atomic separation L_{Cr} . Encircled subsets interact with each other via an effective detuning. (b) On a lattice structure, overlapping pairs have to be used.




FIG. 2. (Color online) Three models consisting of lower order subsets are investigated as simplified models for the $N = 3$ case. The relative deviation $|\rho_{33}^{\text{simpl.}} - \rho_{33}^{\text{exact}}|/\rho_{33}^{\text{exact}}$ from the exact solution is shown for varying detuning Δ and interaction V with the third atom. Models including the exact treatment of pairs give more precise results than the rate equation. The dashed line depicts the models which are suited best for a given V . Parameters: $\Omega_{12} = 3$ MHz, $\Omega_{23} = 2$ MHz, $\gamma_{21} = 6$ MHz, $\gamma_{32} = 25$ kHz, $V_{12} = 2$ MHz.

arise from higher-order processes, which we show to be relevant already at low densities.

We consider a frozen gas [2] of three-level atoms driven by a two-step excitation scheme as shown in Fig. 1. The lower transition between ground state $|1\rangle$ and intermediate state $|2\rangle$ is driven resonantly with Rabi frequency Ω_{12} , and the upper transition between intermediate state $|2\rangle$ and Rydberg state $|3\rangle$ is driven with detuning $\Delta = \omega - \omega_{32}$ and Rabi frequency Ω_{23} .

Before we start with the theoretical analysis, we first illustrate the main idea. We start from a two-atom master equation which yields exact results for two atoms. But there is no unique method to extend this ansatz for pairs to many atoms without using larger atom subsets. To illustrate this, we consider a system of three atoms in which the first two atoms interact strongly with coupling V_{12} . A third atom is moved from large distance ($V := V_{13} = V_{23} = 0$) towards the pair until the three atoms form an equilateral triangle. For the resulting interaction energies V , we compare the exact three-body steady state solution with different simplified models consisting of single atoms or pairs. The result for different detunings Δ is shown in Fig. 2. In the first case (a), we consider the single-atom rate equation model (SARE) [20], in which the interactions are absorbed into effective detunings for each atom individually, and no pairs are formed. The second model (b) accounts for the three possible pairs in the calculation. In addition, each atom in a pair receives an effective detuning due to the interaction with the respective third atom. In the third model we assume that V is smaller than V_{12} , and treat the first two atoms as pair, and the third as individual atom. Again, the mutual interactions between

single atom and pair are included in effective detunings. In Fig. 2 it can be seen that model (a) significantly deviates from the exact treatment at all interaction strengths. Model (b) performs poor at small V , because the overlapping pairs mix the exact and approximate interaction between two atoms. However, (b) is best at high interaction strength. The third ansatz (c) reproduces the exact steady state in the limit of $V \rightarrow 0$ by construction, but fails at large V , as then correlation with atom 3 cannot be neglected. Based on this observation, we construct a model in which the two approaches (b) and (c) are combined, depending on the mutual interactions of the atoms. For this, we introduce a critical interaction (or equivalent: a critical distance) that defines when which model has to be used. This is indicated by the dashed red arrow in Fig. 2. The resulting combination of two approaches provides best results over the whole interaction strength range. The increased complexity of our model due to the inclusion of exact two-atom correlations makes the computation based on established techniques impractically slow. We overcome this by a new method discussed below. Since this approach involves a combination of single-atom and pair descriptions, we call it the hybrid model (HM).

We now proceed with the formal analysis. For a given ensemble of N atoms, we introduce a critical distance L_{Cr} and use it to divide the set of atoms into pairs and single atoms, as schematically shown in Fig. 1. L_{Cr} should be chosen such that many pairs are included to better incorporate two-atom correlations. But on the other hand, overlapping pairs should be avoided as seen in the $N = 3$ case in Fig. 2. Therefore, we use a value which is on the order of the most probable nearest-neighbor distance. The single-particle Hamiltonian in rotating wave approximation reads

$$H^{(i)} = \Omega_{12}(S_{12}^{(i)} + S_{21}^{(i)}) + \Omega_{23}(S_{23}^{(i)} + S_{32}^{(i)}) - \Delta S_{33}^{(i)}. \quad (1)$$

Here $S_{ab}^{(i)}$ denotes the operator $|a\rangle_i \langle b|$ for the i th atom and the Rabi frequencies are assumed to be real. Our hybrid model is based on the full two-body Hamiltonian

$$H^{(i,j)} = H^{(i)} + H^{(j)} + V_{ij} S_{33}^{(i)} S_{33}^{(j)}, \quad (2)$$

with coupling $V_{ij} = C_6/R_{ij}^6$ [1, 2]. Including spontaneous emission as well as dephasing described by the standard Lindblad operator $\mathcal{L}$, the two-atom master equation reads

$$\dot{\rho}^{(i,j)} = -i [H^{(i,j)}, \rho^{(i,j)}] + \mathcal{L}[\rho^{(i,j)}]. \quad (3)$$

In a first attempt, we generalized the SARE approach [20, 21] to solve Eq. (3). But it turned out that the calculation of the rates in the two-atom system is computationally too inefficient. Also, negative rates frequently occur, prohibiting a direct Monte Carlo based solving technique. The latter problem can be overcome




FIG. 3. (Color online) **(a)** Rydberg excitation for a 3D sample of 500 atoms and $C_6 = 50000 \mu\text{m}^6 \text{MHz}$. The hybrid model (dashed lines) and the single atom rate equation (solid lines) yield the same results. **(b)** Rydberg excitation on a 1D lattice for different models with interaction $V_{\text{NN}} = 2.5 \text{ MHz}$ between adjacent atoms. The resonances at $\Delta = V_{\text{NN}}$ and $\Delta = V_{\text{NN}}/2$ correspond to single-atom and two-atom excitations. Other parameters are $\Omega_{12} = 2 \text{ MHz}$, $\Omega_{23} = 1 \text{ MHz}$, $\gamma_{21} = 6 \text{ MHz}$, $\gamma_{32} = 25 \text{ kHz}$, $\Gamma_{32} = \Gamma_{21} = 100 \text{ kHz}$.

by correcting the calculated transition rates to become strictly positive. This eliminates the knowledge of the physical time evolution, but still evolves the system into the correct many-body stationary state [20].

Instead, here, we replace the time-consuming calculation of the rates by an evolution equation based on the readily available steady state solution $\sigma_{ab}^{(\text{SS})}$ of the two-atom populations $\rho_{aa,bb}$ as (see appendix)

$$\dot{\rho}_{aa,bb} = \sum_{c,d} (\sigma_{ab}^{(\text{SS})} - \delta_{ac}\delta_{bd}) \rho_{cc,dd}. \quad (4)$$

By construction, it also propagates the system into the correct stationary state. But as it does not predict the physical intermediate time evolution, we drop all time occurrence in our equations. This allows us to iteratively solve for the many-body steady state with the following three-step Monte Carlo procedure based on the Random Selection Method [10]. First, a pair or a single atom is randomly chosen. Second, the effective detuning for the atom/pair is calculated based on the current configuration of all atoms. With this detuning, the steady state of the atom/pair is calculated. Third, a random number is drawn, and the population of the atom/pair is set in a Monte Carlo step according to the probabilities given by its steady state population distribution. The three steps are repeated until the system converges to the many-body steady state. With our method, several hundred atoms can be simulated easily. Fitting the runtime up to $N = 1000$ results in the scaling law $T \sim N^{1.08}$ and a single realization including 1000 atoms takes about 0.5 seconds on a 3.40 GHz CPU.

In order to verify the validity of the HM we first considered the limit of no pairs, and found results identical to the SARE calculations in [20]. Next, we calculated the Rydberg population probability $\rho_{33} = \sum_i \rho_{33}^{(i)}/N$ and compared the data to established theoretical approaches. In Fig. 3(a), we show results for 3D random samples

of atoms with different gas densities. The suppression of excitation at higher densities can be well understood in terms of the Rydberg blockade. In all cases, our HM agrees perfectly with the SARE [20], indicating the consistency of our method. However, significant deviations occur if the geometry is changed to a 1D lattice as shown in Fig. 1(b). Nearest neighbors interact with $V_{\text{NN}} = 2.5 \text{ MHz}$. The results in Fig. 3(b) show that next to the single-atom excitation peak at $\Delta = 0$, the HM predicts a resonance at $\Delta = V_{\text{NN}}$, and an additional weaker resonance at $\Delta = V_{\text{NN}}/2$. The first condition describes a resonant excitation process for an atom whose neighbor is already excited to the Rydberg state $|3\rangle$, because the laser detuning compensates the interaction energy shift at this point. The second and smaller resonance arises from a 2-photon-process $|22\rangle \rightarrow |33\rangle$ simultaneously exciting two neighboring atoms. The SARE is not capable of producing this second feature because it does not include the exact two-body interaction. To confirm the presence of the 2-photon-resonance we also performed the simulation with the cluster expansion model (CE) used in [14], which also shows this structure. But interestingly, all three approaches significantly deviate already at $\Delta = V_{\text{NN}}$. It should be noted, however, that the lattice geometry is a significant challenge for all models because of overlapping pairs and non-negligible higher-order correlations.

We now turn to our main results on the predictive power of the different models for correlated many-body systems. For this, we calculate the pair correlation function $g^{(2)}(r)$ for a disordered one-dimensional gas. It describes the conditioned probability of having two Rydberg excitations of atoms with distance r [13, 17, 22]. We use the definition [17]:

$$g^{(2)}(r) = \frac{\sum_{i,j}^{(r)} \langle S_{33}^{(i)} S_{33}^{(j)} \rangle}{\rho_{33}^2 \cdot \sum_{i,j}^{(r)} 1}. \quad (5)$$




FIG. 4. (Color online) Pair correlation function $g^{(2)}(r)$ for different laser detunings. **(a)** The models reproduce the Rydberg blockade and the uncorrelated regime. In the intermediate region resonances which indicate an excitation enhancement occur. The CE does not yield the line at $2 \cdot R_1$. The HM data was obtained from averaging 200000 Monte Carlo runs, noise is due to statistics. **(b)** The strong resonance lines are shown in detail and the process $|23\rangle \rightarrow |33\rangle$ at R_1 can be seen. The HM shows $|22\rangle \rightarrow |33\rangle$ at R_2 in addition. Parameters like in Fig. 3, except $C_6 = 900/2\pi \mu\text{m}^6 \text{ MHz}$, $n_{1D} = 0.1 \mu\text{m}^{-1}$.

Here $\sum_{i,j}^{(r)}$ denotes the sum over all pairs of atoms with distance r . For two uncorrelated atoms the pair correlation function is one. Inside the Rydberg blockade regime, where only one excitation can be present, it becomes zero.

Results for $g^{(2)}(r)$ for different values of the laser detuning Δ are shown in Fig. 4. As expected, the blocked and the uncorrelated regimes appear clearly for small and large distances in Fig. 4(a). In between, values $g^{(2)} > 1$ occur for some distances and detunings. These values indicate spatial order, originating from a high probability for multiple excitation at selected distances. Most prominent is the resonance line at about $2 \mu\text{m}$ which can be characterized by the condition $R_1 = (C_6/\Delta)^{1/6}$. Just as in the lattice simulations, it corresponds to the resonant excitation of a second atom when an atom with distance R_1 is already in the Rydberg state. In this case the effective detuning $\Delta_{\text{eff}} = \Delta - C_6/R_1^6$ vanishes. Comparing the results for the HM and CE in Fig. 4(a), it can be seen that the CE does not predict the second peak visible around distances $4 \mu\text{m}$ and for positive detunings in the HM results. This peak occurs if two atoms with distance R_1 are already excited, and a third atom again with distance R_1 to either of the two atoms is excited in addition. The two outermost atoms in this arrangement then have distance $2R_1$. Note that in a higher dimensional geometry this resonance line smears out, because then the distance between the first and the third atom is not necessarily $2R_1$. The CE does not predict this higher-order spatial correlation since triply excited states are not part of the model. In contrast, the SARE method correctly predicts the correlations at $2R_1$.

Next, we analyze the resonance originating from ordering between nearest neighbors, and show a magnified section of Fig. 4(a) in (b). The left panel of (b) shows the results obtained from the SARE, and contains the single resonance at R_1 . But in the result of the HM, another resonance appears. Its condition can be deter-

mined as $R_2 = [C_6/(2\Delta)]^{1/6} \approx 0.89 \cdot R_1$. This resonance again originates from the 2-photon-process $|22\rangle \rightarrow |33\rangle$. Interestingly, in this case, the CE correctly predicts the double resonance, whereas the SARE does not. We note that the calculation of many-body Hamiltonian dynamics in two-level systems also shows, and thus confirms, the discussed resonance lines.

In summary, we presented a hybrid model for the simulation of large ensembles of Rydberg atoms. It is based on an exact two-atom calculation for pairs with distances below a characteristic length scale L_{Cr} , combined with an approximate treatment via effective detunings for more distant atoms. We proposed a method to iteratively solve for the steady state of the HM based on the Random Selection Method without the need for time consuming calculations of transition rates. This way, high predictive power is combined with fast calculation times. We found agreement of the hybrid approach with existing models for simple observables such as the Rydberg excitation probability. But different predictions are found for pair correlation function $g^{(2)}(r)$. The HM predicts structures which both, the CE and the SARE models, can only partially reproduce. We identified the additional structures as originating from multi-atom processes. This not only demonstrates the capability of the HM to characterize higher-order correlations, but also that higher-order correlations cannot be neglected even at low densities.

Financial support by the Heidelberg Center for Quantum Dynamics is gratefully acknowledged.

Appendix

For the propagation of the system into the many-body steady state we use a Monte Carlo algorithm based on the Random Selection Method. In this supplement, technical details of the scheme are described in more detail and a

flowchart of the algorithm is shown in Fig. 5 [23].

Evolution equation

Rate equation based models replace the dynamics of the density matrix ρ by effective equations for the state populations only. For example, for a single atom, if the vector containing all populations is denoted by $\vec{\sigma}$, then

$$\dot{\vec{\sigma}} = A\vec{\sigma},$$

with a matrix A containing rates describing the transitions between the different states. For this reduction to the populations, the dynamics of the coherences is neglected. This in general leads to an approximate description of the dynamics. But the steady state $\vec{\sigma}^{(ss)}$ can be precisely predicted, because there $\dot{\rho}^{(ss)} = 0$ and the dynamics of the coherences is zero by definition. Note that the steady state is unique for quantum optical systems as considered here.

As outlined in the main text, to avoid the problem of negative rates and to significantly speed up our calculation, we instead use the evolution equation (written in component form)

$$\begin{aligned}\dot{\sigma}_a &= \sum_b B_{ab}\sigma_b, \\ B_{ab} &= \sigma_a^{(ss)} - \delta_{ab},\end{aligned}$$




FIG. 5. Schematic algorithm for the hybrid model.

with a tailored evolution matrix B . This matrix is optimized in the sense that it can be calculated much more rapidly than the usual transitions rates in A , as it only depends on the steady state populations $\sigma_a^{(ss)}$. At the same time, it preserves the total probability and evolves the system into the steady state, as $B\vec{\sigma}^{(ss)} = 0$, ensuring consistency with the condition $\dot{\vec{\sigma}}^{(ss)} = B\vec{\sigma}^{(ss)} = 0$. Eq. (4) in the main text is the straightforward generalization of this tailored evolution equation to the two-atom case.

Monte Carlo algorithm

1. *Initialization.* The positions of the N atoms are determined according to the desired density distribution. From the point of view of the algorithm, no restrictions apply to the positions as long as the conditions are such that a perturbative treatment in the order of correlation is meaningful. If the separation of two atoms is smaller than the critical distance L_{Cr} , they are treated as a pair. In total we get N_P pairs and N_A single atoms. Ideally one should have $2N_P + N_A = N$. But in practice, $2N_P + N_A > N$. For example, if three atoms are placed on a line with distance between neighboring atoms smaller than L_{Cr} , then the three atoms count as two pairs, such that $2N_P + N_A = 4 > N = 3$. Next, the initial state of all N atoms is set to the ground state [1].

2. *Choice of atom/pair.* In the first step of the Monte Carlo sequence, the atom or pair is determined whose state will be updated in the sequence. For this, a random integer between 1 and $N_A + N_P$ is drawn which determines the atom or pair which is handled in the current Monte Carlo step. In the following, the indices labeling the atoms are denoted as i in case of a single atom and i_1 and i_2 in case of a pair.

3. *Effective potential.* Next, the effect of all other atoms on the atom/pair chosen in step 2 is calculated. For this, the interaction potential of the atom/pair with all other atoms is summed up and included in an effective detuning. For a single atom i it reads $\Delta_{\text{eff}}^{(i)} = \Delta - \sum_j' V_{ij}$ with the sum $\sum_j'$ running over all other atoms j which are in the Rydberg state [3]. In case of a pair two effective detunings $\Delta_{\text{eff}}^{(i_1)} = \Delta - \sum_{j \neq i_2}' V_{i_1 j}$ and $\Delta_{\text{eff}}^{(i_2)} = \Delta - \sum_{j \neq i_1}' V_{i_2 j}$ have to be calculated. Note that the interaction $V_{i_1 i_2}$ is included exactly in the two-atom description of the pair chosen in step 2.

4. *Steady state.* Next, the steady state of the atom/pair chosen in step 2 under the action of the effective potential calculated in step 3 is determined. For the single atom i , the steady state for the population $\rho_{aa} = \langle S_{aa}^{(i)} \rangle$ is denoted by $\sigma_a^{(ss)}$. For a pair of atoms i_1 and i_2 , the steady state of $\rho_{aa,bb} = \langle S_{aa}^{(i_1)} S_{bb}^{(i_2)} \rangle$ is labeled as $\sigma_{ab}^{(ss)}$. Here, $a, b \in \{1, 2, 3\}$. This step is the most time consuming part of the simulation and should therefore

be highly optimized. By comparing different approaches, we found the following method to be the fastest numerically stable approach. The master equation is linear and can be written as $\dot{\vec{p}} = M \cdot \vec{p}$ where M is a sparse matrix of the size 9×9 or 81×81 for single atoms or pairs, respectively. The fact that the master equation is Hermitian can be exploited to obtain a real matrix M . The problem of finding the steady state is equivalent to find the eigenvector of M with eigenvalue 0. Since all states are coupled, the dimension of the corresponding eigensystem is one and the steady state is unique. M is now decomposed into a product of an orthogonal and an upper triangular matrix in a QR decomposition [24]. Since the matrix M is sparse, Givens Rotations are the best approach to transform M successively into an upper triangular matrix R [24]. R does not have full rank and the last element on the diagonal vanishes. The steady state can then be obtained by back substitution. In practise it is useful to arrange the elements of the vector $\vec{p}$ such that the populations are located at the end, because then the back substitution can be stopped earlier.

5. State update. Next, the state of the atom/pair determined in step 2 is modified according to the population probabilities determined in step 4. For this, a random real number r between 0 and 1 is drawn. In the case of a single atom, the state is changed to $|l\rangle$, where l is the largest integer with $\sum_{k=1}^{l-1} \sigma_k^{(\text{SS})} < r$. For pairs, the procedure is analogous. Note that the new state can be identical to the old state.

6. Loop of the Monte Carlo sequence. The steps 2-5 are now repeated until the system is converged. We have found that for typical parameters, approximately $10 \cdot N$ steps are sufficient to ensure convergence.

7. Calculation of observables. Finally, the observables can be evaluated. Typically, an average over many spatial realizations and Monte Carlo trajectories is necessary to obtain good statistics and smooth results expected from larger ensembles of Rydberg atoms. Note

that throughout the Monte Carlo evolution, each atom is in a definite atomic state at all times, and the diagonal elements of the density matrix only originate from the ensemble averaging. In our work we focus on the evaluation of $g^{(2)}(r)$, and Eq. (5) in the main text shows how to calculate it from the simulation results.

-
- [1] M. Saffman et al., Rev. Mod. Phys. **82**, 2313–2363 (2010).
 - [2] D. Comparat and P. Pillet, J. Opt. Soc. Am. B **27**, A208–A232 (2010).
 - [3] M. D. Lukin et al., Phys. Rev. Lett. **87**, 037901 (2001).
 - [4] D. Jaksch et al., Phys. Rev. Lett. **85**, 2208–2211 (2000).
 - [5] K. Singer et al., Phys. Rev. Lett. **93**, 163001 (2004).
 - [6] E. Urban et al., Nat. Phys. **5**, 110–114 (2009).
 - [7] A. Gaëtan et al., Nat. Phys. **5**, 115–118 (2009).
 - [8] D. Tong et al., Phys. Rev. Lett. **93**, 063001 (2004).
 - [9] H. Weimer et al., Phys. Rev. Lett. **101**, 250601 (2008).
 - [10] A. Chotia et al., New Journal of Physics **10**, 045031 (2008).
 - [11] T. Cubel Liebisch et al., Phys. Rev. Lett. **95**, 253002 (2005).
 - [12] A. Reinhard et al., Phys. Rev. A **78**, 060702 (2008).
 - [13] G. Günter et al., Phys. Rev. Lett. **108**, 013002 (2012).
 - [14] H. Schempp et al., Phys. Rev. Lett. **104**, 173602 (2010).
 - [15] J. D. Pritchard et al., Phys. Rev. Lett. **105**, 193603 (2010).
 - [16] A. Schwarzkopf et al., Phys. Rev. Lett. **107**, 103001 (2011).
 - [17] F. Robicheaux and J. V. Hernández, Phys. Rev. A **72**, 063403 (2005).
 - [18] K. C. Younge et al., Phys. Rev. A **79**, 043420 (2009).
 - [19] B. Olmos et al., Phys. Rev. A **79**, 043419 (2009).
 - [20] C. Ates et al., Phys. Rev. A **83**, 041802 (2011).
 - [21] C. Ates et al., Phys. Rev. A **76**, 013413 (2007).
 - [22] S. Wüster et al., Phys. Rev. A **81**, 023406 (2010).
 - [23] K. P. Heeg, *Models for correlated Rydberg gases*, Diploma thesis (University of Heidelberg, 2011).
 - [24] W. H. Press et al., *Numerical Recipes in FORTRAN: The Art of Scientific Computing*, Vol. 1 (Cambridge University Press, 1992).